



Anti-Human IL-4 Azide Free


PRODUCT SPECIFICATIONS


Catalogue N°	855.030.000 - 200µg / 200µl 855.030.005 - 500µg / 500µl
Target species	Human
Specificity	Recognises both natural and recombinant human IL-4
Clone	B-S4
Application	Immunohistochemistry
Hybridoma	Myeloma X63/AG.8653 x Balb/c spleen cells
Immunisation	Recombinant human IL-4
Quantity	200µg or 500µg (Discovery Size also available please enquire)
Isotype	Mouse IgG1 Kappa light chain
Format	Phosphate-buffered saline. Sterile-filtered through 0.22 µm. Carrier and preservative free
Storage	Stable at +2-8°C for 12 months. For longer storage freeze aliquots.

REFERENCES

Conticello, C. et al., J Immunol.,2004; 172(9): 5467-77. - [Pubmed link](#) 

Stassi, G. et al., Cancer Res.,2003; 63(20): 6784-90. - [Pubmed link](#) 

Todaro, M. et al., Cancer Res.,2006; 66(3): 1491-9. - [Pubmed link](#) 

Venet, F. et al., J Immunol.,2006; 177(9): 6540-7. - [Pubmed link](#) 

BACKGROUND

IL-4 is a lymphokine that co-stimulates the proliferation of activated B- and T-cells, augments the cytotoxic activity of lymphocytes and monocytes and enhances the functional activity of myeloid cells. Produced by mast cells, T-cells and bone marrow stromal cells, IL-4 regulates the differentiation of naive CD4+ cells into helper Th2 cells characterized by their cytokine secretion-profile that includes secretion of IL-4, IL-5, IL-6, IL-10 and IL-13 which favour a humoral immune response. In addition, IL-4 can inhibit the proliferation of TNF, IL-1 and IL-6 by macrophages.

Discover Diaclone

With over 30 years experience and extensive expertise, we are committed to providing excellence in Monoclonal Antibody and Immunoassay development.

The expanding range of Diaclone Immunology products is specifically designed to advance research applications.

Our experience and expertise coupled to the diversity and quality of our product range makes Diaclone a clear choice to :

Fast Track Your Research

Diaclone products



6 rue Dr Jean-François-Xavier Girod
BP 1985
25020 Besançon cedex
France

Tel : +33 (0)3 81 41 38 38
Fax : +33 (0)3 81 41 36 36

Email : Info@diaclone.com
www.diaclone.com

IL-4 induces B-cell class switching to IgE and IgG1 isotypes, and up regulates MHC Class II production and CD23 expression.

IL-4 is a 15kDa globular glycoprotein containing 129 amino acid residues. The non glycosylated form of the protein is fully biologically active.

Version 9 - 05.21

For research use only